



CLAIMS

1. A dye mixture comprising,
as component (A), a reactive dye of the formula (A)



wherein each of Y^1 and Y^2 , independently, is a vinyl group or a group of the formula $-CH_2CH_2Q$ in which Q is a leaving group removable under alkaline conditions to provide a vinyl group; and,

as component (B), at least one reactive dye selected from
(I) a monoazopyrazole dye of the formula



wherein: R^1 is an aryl group selected from phenyl and naphthyl groups optionally having at least one substituent thereon, the substituents, or each substituent independently, being selected from a sulphonic acid group and a salt thereof, a

C₁₋₄ alkyl group, a C₁₋₄alkoxy group, a hydroxy group, a carboxyl group, a chlorine atom, a vinyl sulphonyl group and a group SO₂CH₂CH₂Q¹ in which Q¹ is a leaving group removable under alkaline conditions to provide a vinyl sulphonyl group;

R² is a phenyl or naphthyl group, optionally having at least one substituent thereon, the substituent, or each substituent, independently, being selected from a sulphonic acid group and a salt thereof, a C₁₋₄alkyl group, a C₁₋₄alkoxy group, a hydroxy group, a chlorine atom, a vinyl sulphonyl group, a group SO₂CH₂CH₂Q² in which Q² is a leaving group removable under alkaline conditions to provide a vinyl sulphonyl group, a group Het and a group L⁵-Het, where Het is an optionally substituted aromatic heterocyclic reactive or non-reactive group or a reactive or non-reactive group having an aliphatic chain and L⁵ is as defined below; and

R⁶ is a methyl group, an amide group or a carboxyl group or a salt thereof;


each of R⁴⁰ and R⁴¹, independently, is an aryl group selected from phenyl and naphthyl groups, each of which, independently, is optionally substituted by a vinylsulphonyl group, a group SO₂CH₂CH₂Q¹ in which Q¹ is a leaving group removable under alkaline conditions to provide a vinylsulphonyl group; or the group Het, where Het is as defined above;

at least one of R^1 , R^2 , R^{40} and R^{41} being, or having thereon at least one substituent which is, reactive;

each of L^1 , L^2 and L^5 independently is a linking group selected from $N(R^{20})$, in which R^{20} is hydrogen or C_{1-4} alkyl; $C(=O)$; $C(=O)-O$; $S(=O)_2$; $S(=O)-NH$; $C(=O)-NH$; and $NHC(=O)NH$; and


each of x , y and z , independently, is zero or 1; and, when the group R^1 is substituted by a hydroxy group ortho to the azo group, a metallized derivative thereof;

(II) a monoazo or disazo dye of the formula



wherein: Het is an optionally substituted aromatic heterocyclic reactive or non-reactive group or a reactive or non-reactive group having an aliphatic chain;

R^3 or each R^3 , independently, is a chlorine atom, a methyl group, a methoxy group, a sulphonic acid group or a salt thereof, or is an amino group of the formula



in which each of R^{18} and R^{19} , independently, is hydrogen, chloro, methyl, (C_{1-4} alkyl)carbonyl, aminocarbonyl, vinylsulphonyl or a group $SO_2CH_2CH_2Q^1$, in which Q^1 is as defined above;

R^4 , or each R^4 independently, is hydrogen, a sulphonic acid group or a salt thereof, a C_{1-8} alkyl group, a C_{1-4} alkoxy group, a vinyl sulphonyl group or a group $SO_2CH_2CH_2Q^2$ in which Q^2 is a leaving group removable under alkaline conditions to provide a vinyl sulphonyl group, which C_{1-4} alkyl group or C_{1-4} alkyl moiety of the C_{1-4} alkoxy group is optionally interrupted by an oxygen atom to provide an ether group and is optionally substituted by a vinyl sulphonyl group or a group $SO_2CH_2CH_2Q^2$, in which Q^2 is as defined above; or R^4 (or when q is 2, each R^4 independently) is a phenyl group optionally substituted by at least one sulphonic acid group or a salt thereof or at least one group Het, where Het is as defined above; or R^4 is a group Het, where Het is as defined above;

R^5 is an aryl group selected from phenyl and naphthyl groups each optionally substituted by at least one sulphonic acid group or a salt thereof or at least one group Het, as defined above;

R^{20} is a hydrogen atom or a C_{1-4} alkyl group;

R^{21} is a hydrogen atom, a C_{1-4} alkyl group, a sulphonic acid- C_{1-4} alkyl group, a chloroalkylsulphonyl- C_{1-4} alkyl group or a group Het, where Het is as defined above;

m is 1 or 2;

p is zero, 1 or 2;

q is zero, 1 or 2;


each of w_1 and w_2 is zero or 1; and

w_3 is 1, 2 or 3; and

when p is zero, q is zero;

at least one of R^4 , R^5 , R^{18} , R^{19} , R^{21} and Het being, or having thereon at least one substituent which is, reactive.

(III) a monoazopyridone dye of the formula



wherein: each of R^{40} , R^{41} , L^1 and L^2 is as defined above;

R^7 is optionally present, and is a cyano group or the group $-CH_2SO_3H$ or the group $-C(=O)NH_2$;

each of R^{24} and R^{25} , independently, is a hydrogen atom, a C_{1-4} alkyl group, a sulphonyl- C_{1-4} alkyl group, or a carboxyl group;

n is 1 or 2;


r is zero or 1; and

each of s_1 and s_2 is zero or 1; and when R^{40} is a phenyl or naphthyl group s_2 is 1;

at least one of R^{40} and R^{41} being, or having thereon

at least one substituent which is, reactive;

(IV) a monoazonaphthyl dye of the formula



wherein: each of R^1 , R^{40} , R^{41} , L^1 , L^2 , x , y and z is as defined above;

R^9 is $\text{CH}_3(\text{C}=\text{O})-$, Het (as defined above) or an aryl group selected from phenyl and naphthyl, which Het or aryl group is optionally substituted by at least one substituent, the or each substituent, independently, being selected from a sulphonic acid group and a salt thereof, a C_{1-4} alkyl group, a C_{1-4} alkoxy group, a hydroxy group, an amino group optionally substituted by at least one methyl or sulphato group, a vinyl sulphonyl group and a group $\text{SO}_2\text{CH}_2\text{CH}_2\text{Q}^1$ in which Q^1 is as defined above;

R^{35} is a C_{1-4} alkyl or C_{2-4} alkenyl group, which C_{1-4} alkyl or C_{2-4} alkenyl group is optionally substituted by at least one halogen atom, a sulphonic acid group or salt thereof, a chloroalkylsulphonyl group, a vinylsulphonyl group or $-\text{SO}_2\text{CH}_2\text{CH}_2\text{Q}^1$, where Q^1 is as defined above and which C_{1-4} alkyl or C_{2-4} alkenyl group optionally additionally contains at least one oxygen or sulphur atom in the chain thereof; the

group Het (as defined above); or an aryl group selected from phenyl and naphthyl, which Het or aryl group is optionally substituted by at least one substituent, the substituent or each substituent independently, being selected from a sulphonic acid group and a salt thereof, a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group, a halogen atom, a hydroxy group, an amino group optionally substituted by at least one methyl or sulphato group, a vinylsulphonyl group, a vinylsulphonyloxyethyl group and a group SO₂CH₂CH₂Q¹ in which Q¹ is as defined above;

R⁵⁰ is a vinylsulphonyl group or a group SO₂CH₂CH₂Q¹ (in which Q¹ is as defined above);

at least one of R¹, R⁹, R³⁵, R⁴⁰, R⁴¹ and R⁵⁰ is, or has thereon at least one substituent which is, reactive;

L³ is a linking group selected from N(R²⁰), in which R²⁰ is as defined above; CO; COO; NHCO; NHCONH; SO₂NH and SO₂;

L⁴ is a linking group selected from N(R²¹), in which R²¹ is as defined above, CO, COO, NHCO, NHCONH, SO₂NH and SO₂;

Z is hydroxy, amino or methylamino;

a is zero or 1-4;

b is zero or 1-3;


c is zero or 1;

d is zero, 1 or 2;

e is zero or 1; and

when each of R^1 and Z provides a hydroxyl group ortho to the azo group, a metallized derivative thereof;

(V) a disazoaminonaphthyl dye of the formula



wherein: each of R^1 , R^{40} , R^{41} , L^1 , L^2 , x , y and z is as defined above;

each of V and W , independently, is NH_2 or OH ;

R^{14} is an aryl group selected from phenyl and naphthyl groups optionally having at least one substituent thereon, the substituents, or each substituent independently, being selected from a sulphonic acid group and a salt thereof, a C_{1-4} alkyl group, a C_{1-4} alkoxy group, a hydroxy group, a vinyl sulphonyl group, a group $\text{SO}_2\text{CH}_2\text{CH}_2\text{Q}^1$ in which Q^1 is a leaving group removable under alkaline conditions to provide a vinyl sulphonyl group;

each of R^{51} and R^{52} independently is an aryl group


selected from phenyl and naphthyl groups each of which is optionally substituted by a vinyl sulphonyl group, a group $\text{SO}_2\text{CH}_2\text{CH}_2\text{Q}^1$ in which Q^1 is a leaving group removable under alkaline conditions to provide a vinyl sulphonyl group, or the group Het^3 , where Het^3 is an optionally substituted aromatic heterocyclic reactive group or a reactive group having an aliphatic chain;

each of L^3 and L^4 , independently, is a linking group selected from $\text{N}(\text{R}^{20})$, in which R^{20} is hydrogen or C_{1-4} alkyl; $\text{C}(=\text{O})$; $\text{C}(=\text{O})-\text{O}$; $\text{S}(=\text{O})_2$; $\text{S}(=\text{O})-\text{NH}$; $\text{C}(=\text{O})-\text{NH}$; and $\text{NHC}(=\text{O})\text{NH}$;

each of f , g and h , independently is zero or 1; and

at least one of R^{14} , R^{40} , R^{41} , R^{51} and R^{52} is, or has thereon at least one substituent which is, reactive.

(VI) a formazan dye of the formula



wherein: each of R^{16} and R^{17} independently of one another,


each R^{16} independently of one another and each R^{17} independently of one another, is a sulphonic acid group or a salt thereof, a vinyl sulphonyl group or a group $SO_2CH_2CH_2Q^1$ in which Q^1 is a leaving group removable under alkaline conditions to provide a vinyl sulphonyl group;

each of Het^1 and Het^2 , independently, is an optionally substituted aromatic heterocyclic reactive or non-reactive group or a reactive or non-reactive group having an aliphatic chain; and

each of R^{60} and R^{61} , independently, is an aryl group selected from phenyl and naphthyl groups each of which is optionally substituted by a sulphonic acid group or a salt thereof, a vinylsulphonyl group or a group $SO_2CH_2CH_2Q^1$ in which Q^1 is as defined above;

X^1 is a sulphonic acid group or a salt thereof or halogen; each of i , j and v , independently, is zero, 1 or 2; and each of k , l , t and u , independently, is zero or 1; and


at least one of R^{16} , R^{17} , R^{50} , R^{51} , Het^1 and Het^2 is, or has thereon at least one substituent which is, reactive;



wherein: each of R^1 , R^2 , R^3 , R^4 , R^{20} , R^{21} , R^{40} , L^1 , Het, x, p, q, w_1 , w_2 and w_3 is as defined above; and

at least one of R^1 , R^2 , R^3 , R^4 , R^{21} , R^{40} and Het is, or has thereon at least one substituent which is, reactive;

(VIII) a disazo dye of the formula



wherein R^{55} is an aryl group selected from phenyl and naphthyl groups;

R^{56} is a sulphonic acid group or a salt thereof or a reactive group selected from a vinyl sulphonyl group and a group $SO_2CH_2CH_2Q^1$ in which Q^1 is a leaving group removable under alkaline conditions to provide a vinyl sulphonyl group;

R^{57} is an amino group or a group NHR^A in which R^A is a C_{1-4} alkyl group;

R^{58} is a sulphonic acid group or a salt thereof;

R^{59} is a sulphonic acid group or a salt thereof, a reactive group selected from a vinyl sulphonyl group and a group $SO_2CH_2CH_2Q^2$ in which Q^2 is a leaving group removable under alkaline conditions to provide a vinyl sulphonyl group; or the

group R^{59} is a group Het or a group L^{10} -Het, where Het is an optionally substituted aromatic heterocyclic reactive or non-reactive group;

R^{65} is a ureido group or a group $HNC(=O)R^B$ in which R^B is a C_{1-4} alkyl group;

L^{10} is a linking group selected from $N(R^{20})$, in which R^{20} is hydrogen or C_{1-4} alkyl; $C(=O)$; $C(=O)-O$; $S(=O)_2$; $S(=O)-NH$; $C(=O)-NH$; and $NHC(=O)NH$;

y_1 is zero, 1, 2 or 3;


y_2 is zero, 1 or 2;

y_3 is zero or 1; and


y_4 is zero or 1; and

at least one of R^{56} and R^{59} is a reactive group.


(IX) a dye of the formula



or



(X) a metal phthalocyanine dye of the formula



wherein: MPc is a metallophthalocyanine chromophore;

each of R^{70} and R^{71} , independently, is hydrogen or C_{1-4} alkyl;

B is a hydrocarbon bridging group;

Het is a reactive heterocyclic group;

each of x_1 , x_2 and x_3 is a respective average value;


$$x_1 + x_2 + x_3 = 4;$$

x_1 is at least 1

x_2 is zero or 1; and

x_3 is at least 1; and

(XI) a triphenodioxazine dye of the formula (XI) (or a salt thereof)



wherein: each of B^1 and B^2 , independently, is a hydrocarbon bridging group;

U^1 is H or SO_3H ; and


each of T^1 and T^2 , independently, is halo, C_{1-4} alkyl, or C_{1-4} alkoxy;

each of R^{80} and R^{81} is a phenyl group substituted by at least one sulphonic acid group or a salt thereof;

each of Het^1 and Het^2 is as defined above; and

at least one of Het^1 and Het^2 is a reactive group.

2. A dye mixture according to claim 1, wherein, in the dye of the formula (A), at least one of Y^1 and Y^2 is the group $-CH_2CH_2Q$ and Q is selected from chlorine, bromine, C_{1-4} -alkylsulfonyl, phenylsulfonyl, OSO_3H , SSO_3H , $OP(O)(OH)_2$, C_{1-4} -alkylsulfonyloxy, phenylsulfonyloxy, $(C_{1-4}$ alkyl) carbonyloxy, $(C_{1-4}$ dialkyl) amino or a radical of the formula



where Z^9 , Z^{10} and Z^{11} are identical or different and are each, independently of one another, C_{1-4} alkyl or benzyl and An^\ominus is in each case one equivalent of an anion.

3. A dye mixture according to claim 2, in which, in the reactive dye of the formula (A), each of Y^1 and Y^2 is the group $HO_2SOCH_2CH_2$.


4. A dye mixture according to any preceding claim, wherein the reactive dye (B) contains a group Het, where Het is an optionally substituted aromatic heterocyclic reactive group derived from a halogen-substituted heterocyclic compound selected from 1,3,5-triazine, quinoxaline, phthalazine, pyrimidine, pyridazine and 2-(C₁₋₄ alkylsulphonyl) benzothiazole.

5. A dye mixture according to claim 4, wherein the aromatic heterocyclic reactive group is substituted and at least one substituent is a halogen atom.

6. A dye mixture according to any one of claims 1 to 3, wherein the reactive dye (B) contains a group Het, where Het is a reactive group having an aliphatic chain and selected from acryloyl, mono-, di- or trichloroacryloyl, mono-, di- or tri- bromoacryloyl, -CO-CCl=CH-COOH, -CO-CH=CCl-COOH, 2-chloropropionyl, 1,2-dichloropropionyl, 1,2-dibromopropionyl, 3-phenylsulfonylpropionyl, 3-methylsulfonylpropionyl, 2-sulfatoethylaminosulfonyl, 2-chloro-2,3,3-trifluorocyclobutylcarbonyl, 2,2,3,3-tetrafluorocyclobutylcarbonyl, 2,2,3,3-tetrafluorocyclobutylsulfonyl, 2-(2,2,3,3-tetrafluorocyclobutyl)acryloyl, 1- or 2-alkyl- or 1- or 2-arylsulfonylacryloyl, or a radical of the formula SO₂-Y³, SO₂NH-Y³, CONH-L⁶-SO₂-Y³ or NHCONH-L⁶-SO₂-Y³ where L⁶ is C₁-C₄-alkylene or phenylene and Y³ is a vinyl group or a group of the

formula $\text{CH}_2\text{CH}_2\text{Q}^3$ in which Q^3 is a leaving group removable under alkaline conditions to provide a vinyl group.

7. A dye mixture according to any preceding claim, wherein the reactive dye (B) has at least one of the groups Q^1 and Q^2 therein and the or each of groups Q^1 and Q^2 independently is selected from chlorine, bromine, C_{1-4} -alkylsulfonyl, phenylsulfonyl, OSO_3H , SSO_3H , $\text{OP}(\text{O})(\text{OH})_2$, C_{1-4} -alkylsulfonyloxy, phenylsulfonyloxy, $(\text{C}_{1-4}$ alkyl)carbonyloxy, $(\text{C}_{1-4}$ dialkyl)amino or a radical of the formula



where Z^9 , Z^{10} and Z^{11} are identical or different and are each, independently of one another, C_{1-4} -alkyl or benzyl and An^- is in each case one equivalent of an anion.

8. A dye mixture according to any preceding claim, wherein the reactive dye (B) is a monoazopyrazole dye of the formula (I), given and defined in claim 1.

9. A dye mixture according to claim 8, wherein, in the reactive dye (B) of the formula (I), at least one of the group $[(\text{R}^{41}-\text{L}^2)_z(\text{R}^{40}-\text{L}^1)_y]\text{R}^1$ and the group R^2 has a group $-\text{SO}_2\text{CH}_2\text{SO}_2\text{Q}^2$ substituted thereon, Q^2 being as defined in claim 1.

10. A dye mixture according to claim 8 or claim 9, wherein, in the reactive dye (B) of the formula (I), x is zero and the group R^1 is substituted by at least one of a sulphonic acid group and the group $-SO_2CH_2SO_2Q^1$, wherein Q^1 is as defined in claim 1.

11. A dye mixture according to claim 8 or claim 9, wherein, in the reactive dye (B) of the formula (I), each of x, y and z is 1, each of L^1 and L^2 is NH, R^{40} is the group Het, where Het is a triazine ring substituted by a halogen atom and R^{41} is an optionally substituted phenyl group.


12. A dye mixture according to claim 8 or claim 9, wherein, in the reactive dye (B) of the formula (I), each of x and z is 1, y is zero, L^2 is NH and R^{41} is the group Het, where Het is a difluorochloropyrimidinyl group.

13. A dye mixture according to claim 8 or claim 9, wherein, in the reactive dye (B) of the formula (I), each of x and z is 1, y is zero, L^2 is CONH (in which the nitrogen atom is attached to the group R^1 and the carbon to the group R^{41}) and R^{41} is the group Het, where Het is a 2,3-dichloroquinoxaline group.

14. A dye mixture according to any one of claims 8 to 13, wherein, in the reactive dye (B) of the formula (I), R^2 is a

phenyl or naphthyl group substituted by at least one of a sulphonic acid group and the group $-\text{SO}_2\text{CH}_2\text{SO}_2\text{Q}^2$, where Q^2 is as defined in claim 1.

15. A dye mixture according to any one of claims 8 to 13, wherein, in the reactive dye (B) of the formula (I), R^2 is a phenyl or naphthyl group substituted at least by the group NH-Het , where Het is a triazine ring substituted by a halogen atom and optionally substituted by the group



16. A dye mixture according to any one of claims 8 to 15, wherein, in the reactive dye (B) of the formula (I), the group R^1 is substituted by a hydroxy group ortho to the azo group and the reactive dye (B) is in the form of a copper complex.

17. A dye mixture according to any one of claims 1 to 7, wherein the reactive dye (B) is monoazo or disazo dye of the formula (II), given and defined in claim 1.

18. A dye mixture according to claim 17, wherein, in the reactive dye (B) of the formula (II), the group R^5 is an aryl group selected from phenyl and naphthyl groups each substituted by at least one sulphonic acid group or a salt thereof or by a

group Het, wherein Het is a vinylsulphonyl group or a group $\text{SO}_2\text{CH}_2\text{CH}_2\text{Q}^1$ in which Q^1 is as defined in claim 1.

19. A dye mixture according to claim 16 or claim 18, wherein, in the reactive dye (B) of the formula (II), m is 1, which dye is a monoazo dye.

20. A dye mixture according to claim 19, wherein, in the reactive dye (B) of the formula (II), each of w_1 , w_2 , p and q is zero, w_3 is at least 2, at least one of the groups R^3 is a sulphonic acid group and the group R^5 is an aryl group selected from phenyl and naphthyl groups each substituted by a vinylsulphonyl group or a group $-\text{SO}_2\text{CH}_2\text{CH}_2\text{Q}^1$, where Q^1 is as defined in claim 1.


21. A dye mixture according to claim 19, wherein, in the reactive dye (B) of the formula (II), each of w_1 and w_2 is zero, each of p and q is 1, R^{21} is hydrogen and R^4 is selected from a triazine ring substituted by at least one halogen atom and optionally additionally substituted by an amino group; and a pyrimidine group substituted by at least one halogen atom and optionally additionally substituted by a methyl group.

22. A dye mixture according to claim 19, wherein, in the reactive dye (B) of the formula (II), each of w_1 , w_2 , p and q is 1, R^{20} is hydrogen, Het is a triazine ring substituted by a


halogen atom, R^{21} is hydrogen and R^4 is selected from a phenyl group or a group $-SO_2CH_2CH_2Q^1$, where Q^1 is as defined in claim 1, and a straight or branched C_{2-4} alkylene chain substituted by a substituent selected from a hydroxyl group; a sulphonic acid group or salt thereof; a vinylsulphonyl group, a group $-SO_2CH_2CH_2Q^2$, where Q^2 is as defined in claim 1; and a pyrimidinylamino group in which the pyrimidinyl group is substituted by at least one halogen atom and optionally additionally by a methyl group; and which straight or branched C_{2-4} alkylene group optionally contains a hetero atom selected from O, S and N(H).

23. A dye mixture according to claim 19, wherein, in the reactive dye (B) of the formula (II), each of w_1 and w_2 is zero, each of p and q is 1 and each of R^{21} and R^4 is the group OSO_3H .

24. A dye mixture according to claim 19, wherein, in the reactive dye (B) of the formula (II), w_1 is zero, w_2 is 1, p is 1, q is 1, R^{21} is hydrogen, R^4 is a pyrimidinyl group substituted by at least one halogen atom and optionally additionally substituted by a methyl group and Het is a non-reactive heterocyclic group of the formula



25. A dye mixture according to claim 19, which contains a dye of the formula (44)



26. A dye mixture according to any one of claims 1 to 7, wherein the reactive dye (B) is a monoazopyridone dye of the formula (III), given and defined in claim 1.

27. A dye according to claim 26, wherein, in the reactive dye (B) of the formula (III), each of r , s_1 and s_2 is 1, each of L^1 and L^2 is NH, R^{40} is a triazine ring substituted by a halogen atom and R^{41} is selected from a phenyl group substituted by at least one of a chlorine atom, a sulphonic acid group or a salt thereof, a vinylsulphonyl group or a group $-SO_2CH_2CH_2Q^1$, where Q^1 is as defined in claim 1; and a straight or branched chain C_{2-4} alkylene group optionally containing at least one oxygen atom and substituted by a vinylsulphonyl group or a group $-SO_2CH_2CH_2Q^1$, where Q^1 is as defined in claim 1.

28. A dye mixture according to claim 26, wherein, in the reactive dye (B) of the formula (III), each of r and s_1 is 1, s_2 is zero, L^2 is NH, R^{41} is a pyrimidinyl group substituted by at least one halogen atom and optionally additionally substituted by a methyl group and R^{40} is a naphtho-(1,2-d)-1,2,3-triazole substituted by at least one sulphonic acid group or salt thereof.

29. A dye mixture according to claim 26, wherein, in the reactive dye (B) of the formula (III), each of r_1 , s_1 and s_2 is 1, L^1 is CONH (in which the carbon atom is attached to the group R^{40}), L^2 is NH, R^{40} is an optionally substituted phenyl group and R^{41} is a pyrimidinyl group substituted by at least one halogen atom and optionally additionally substituted by a methyl group.

30. A dye mixture according to any one of claims 1 to 7, wherein the reactive dye (B) is a monoazonaphthyl dye of the formula (IV), given and defined in claim 1.

31. A dye mixture according to claim 30, wherein, in the reactive dye (B) of the formula (IV), x is zero, d is 1 and a or b is 1.

32. A dye mixture according to claim 31, wherein e is zero.

33. A dye mixture according to claim 30, wherein, in the reactive dye (B) of the formula, x is 1, y is zero, z is 1, L^2 is NH and R^{41} is a pyrimidinyl group substituted by at least one halogen atom and optionally additionally substituted by a methyl group.

34. A dye mixture according to claim 30, wherein, in the reactive dye (B) of the formula (IV), x is 1, y is zero, z is 1, L² is CONH (with the carbon atom attached to the group R¹ and the nitrogen atom to the group R⁴¹) and R⁴¹ is a phenyl group substituted by a vinylsulphonyl group or a group -SO₂CH₂CH₂Q¹, where Q¹ is as defined in claim 1.

35. A dye mixture according to claim 30, wherein, in the reactive dye (B) of the formula (IV) each of x, y and z is 1, each of L¹ and L² is NH, R⁴⁰ is a triazine ring substituted by a halogen atom and R⁴¹ is an aromatic group selected from phenyl and naphthyl groups each substituted by at least one of a sulphonic acid group or a salt thereof, a vinylsulphonyl group and the group -SO₂CH₂CH₂Q¹, where Q¹ is as defined in claim 1.

36. A dye mixture according to any one of claims 33 to 35, wherein e is zero.

37. A dye mixture according to claim 30, wherein, in the reactive dye (B) of the formula (IV), e is 1, c is zero, L³ is CONH (where either the nitrogen or carbon atom thereof is attached to the group R⁹) and R⁹ is a methyl group.

38. A dye mixture according to claim 30, wherein, in the reactive dye (B) of the formula (IV), e is 1, c is zero, L³ is CONH (where either the nitrogen or carbon atom thereof is

attached to the group R^9) and R^9 is a phenyl group optionally substituted by at least one substituent, the or each substituent being selected from methoxy, carboxyethyl, sulphoethyl, carboxyethenyl, 1,2-dibromoalkyl, chloroethylsulphonyl, vinylsulphonyl, a group of the formula $-SO_2CH_2CH_2Q^1$, where Q^1 is as defined in claim 1 and a 2,3-dichloroquinoxaline group.

39. A dye mixture according to claim 30, wherein, in the reactive dye (B) of the formula (IV), e is 1, c is 1, L^3 is CONH (where either the nitrogen or carbon atom thereof is attached to the group R^9), R^9 is a phenyl group substituted at least by the group L^4-R^{35} , where L^4 is NH and R^{35} is a vinyl group optionally substituted by a halogen atom or a pyrimidinyl group substituted by a halogen atom and optionally additionally substituted by a methyl group.

40. A dye mixture according to claim 30, wherein, in the reactive dye (B) of the formula (IV), e is 1, c is zero, L^3 is $N(R^{20})$, where R^{20} is as defined in claim 1 and R^9 is a vinylsulphonyl group, a group $-SO_2CH_2CH_2Q^1$ wherein Q^1 is as defined in claim 1, or a pyrimidinyl group substituted by a halogen atom and optionally additionally substituted by a methyl group.

41. A dye mixture according to claim 40, wherein, in the group $N(R^{20})$, R^{20} is a halogen atom or a methyl group.

42. A dye mixture according to claim 30, wherein, in the reactive dye (B) of the formula (IV), e is 1, c is zero, L^3 is $N(R^{20})$, where R^{20} is as defined in claim 1 and R^9 is a triazine ring substituted by at least one halogen atom.

43. A dye mixture according to claim 42, wherein the triazine ring is substituted by two halogen atoms.


44. A dye mixture according to claim 42, wherein the triazine ring is substituted by one halogen atom and a morpholinyl group.

45. A dye mixture according to claim 30, wherein, in the reactive dye (B) of the formula (IV), e is 1, c is 1, L^3 is $N(R^{20})$, where R^{20} is as defined in claim 1, L^4 is the group R^{21} , where R^{21} is as defined in claim 1 and R^{35} is selected from a phenyl group optionally substituted by a sulphonic acid group or salt thereof, a halogen atom, a vinylsulphonyloxyalkyl group, a vinylsulphonyl group or the group $-SO_2CH_2CH_2Q^1$, where Q^1 is as defined in claim 1; and a C_{1-4} alkyl group optionally substituted by a vinylsulphonyl group, the group $-SO_2CH_2CH_2Q^1$, where Q^1 is as defined in claim 1, a sulphonic acid group or a salt thereof or a chloroalkylsulphonyl group, which C_{1-4} alkyl group optionally additionally contains at least one oxygen or sulphur atom in the chain thereof.

46. A dye mixture according to any one of claims 37 to 45, wherein x is zero.

47. A dye mixture according to any one of claims 30 to 46, wherein, in the reactive dye (B) of the formula (IV), each of the groups R^1 and the naphthalene nucleus is substituted by a respective hydroxyl group ortho to the azo group and the reactive dye (B) is in the form of a copper complex thereof.

48. A dye mixture according to claim 30, which contains a dye of the formula (53)



49. A dye mixture according to any one of claims 1 to 7, wherein the reactive dye (B) is a disazo dye of the formula (V), given and defined in claim 1.

50. A dye mixture according to claim 49, wherein, in the reactive dye (B) of the formula (V), V is amino, W is hydroxy, each of R^1 and R^4 is a phenyl group substituted by at least one substituent, the or each substituent independently being selected from a sulphonic acid group or a salt thereof, a

vinylsulphonyl group and a group $\text{SO}_2\text{CH}_2\text{CH}_2\text{Q}^1$, where Q^1 is as defined in claim 1 and R^{14} is optionally additionally substituted by at least one methoxy group.

51. A dye mixture according to claim 49 or claim 50, wherein, in the reactive dye (B) of the formula (V), each of x and h is zero.

52. A dye mixture according to claim 49 or claim 50, wherein, in the reactive dye (B) of the formula (V), at least one of x and h is 1.

53. A dye mixture according to claim 52, wherein, each of x and h is 1.

54. A dye mixture according to claim 50, wherein each of x, y and z is 1, each of L^1 and L^2 is NH , R^{40} is a triazine ring substituted by a halogen atom and R^{41} is a phenyl group substituted by at least one substituent, the or each substituent independently being selected from a sulphonic acid group or a salt thereof, a vinylsulphonyl group and a group $\text{SO}_2\text{CH}_2\text{CH}_2\text{Q}^1$, where Q^1 is as defined in claim 1.

55. A dye mixture according to any one of claims 50 and 52 to 54, wherein each of f, g and h is 1, each of L^3 and L^4 is NH , R^{51} is a triazine ring substituted by a halogen atom and R^{52} is

a phenyl group substituted by at least one substituent the or each substituent independently being selected from a halogen atom, a sulphonic acid group or a salt thereof, a vinylsulphonyl group and a group $\text{SO}_2\text{CH}_2\text{CH}_2\text{Q}^1$, where Q^1 is as defined in claim 1.

56. A dye mixture according to claim 49, wherein, in the reactive dye (B) of the formula (V), V is amino, W is hydroxy, h is zero and R^{14} is a naphthalene group substituted by at least one sulphonic acid group.

57. A dye mixture according to claim 56, wherein each of x, y and z is 1, R^1 is a phenyl group optionally substituted by a sulphonic acid group or a salt thereof, each of L^1 and L^2 is NH, R^{40} is a triazine ring substituted by a halogen atom and R^{41} is a phenyl group substituted by at least one substituent, the or each substituent independently being selected from a sulphonic acid group or a salt thereof, a vinylsulphonyl group and a group $\text{SO}_2\text{CH}_2\text{CH}_2\text{Q}^1$, where Q^1 is as defined in claim 1.

58. A dye mixture according to claim 49, wherein, in the reactive dye (B) of the formula (V), V is amino, W is hydroxy, h is 1, f is zero, g is 1, L^4 is NH and R^{52} is the group Het^3 , where Het^3 is a substituted aromatic heterocyclic group.

59. A dye mixture according to claim 58, wherein the group Het^3 is a pyrimidinyl group substituted by at least one halogen atom and optionally additionally substituted by a methyl group.


60. A dye mixture according to claim 59, wherein x is zero and R^1 is a phenyl group substituted by a vinylsulphonyl group or $SO_2CH_2CH_2Q^1$, where Q^1 is as defined in claim 1.

61. A dye mixture according to claim 49, wherein, in the reactive dye (B) of the formula (V), V is amino, W is hydroxy, h is 1, f is zero, g is 1, L^4 is $NHCO$ (where either the nitrogen or carbon atom thereof is attached to the group R^{14}) and R^{52} is the group Het^3 , where Het^3 is a reactive group having an aliphatic chain.


62. A dye mixture according to claim 61, wherein the group Het^3 is a vinylsulphonyl group or $SO_2CH_2CH_2Q^1$, where Q^1 is as defined in claim 1.

63. A dye mixture according to claim 62, wherein x is zero and R^1 is a phenyl group substituted by a vinylsulphonyl group or $SO_2CH_2CH_2Q^1$, where Q^1 is as defined in claim 1.


64. A dye mixture according to claim 49, which contains a dye of the formula (105)




65. A dye mixture according to claim 64, which additionally contains a dye of the formula (44)



66. A dye mixture according to claim 49, which contains a dye of the formula (163)



67. A dye mixture according to claim 49, which contains a dye of the formula (167)



68. A dye mixture according to claim 66 or 67, which contains each of the dyes of the formulae (163) and (167).

69. A dye mixture according to any one of claims 1 to 7, wherein the reactive dye (B) is a formazan dye of the formula (VI) given and defined in claim 1.

70. A dye mixture according to claim 69, wherein, in the reactive dye (B) of the formula (VI), each of u and v is zero, i is 1, R^{16} is a sulphonic acid group or a salt thereof, j is 2 and one R^{17} is a sulphonic acid group or a salt thereof and the other R^{17} is a vinylsulphonyl group or $SO_2CH_2CH_2Q^1$, where Q^1 is as defined in claim 1.

71. A dye mixture according to claim 69, wherein, in the reactive dye (B) of the formula (VI), v is zero, i is 1, R^{16} is a sulphonic acid group or a salt thereof, j is 1, R^{17} is a sulphonic acid group, u is 1 and Het^1 is a triazine ring substituted by a halogen atom.

72. A dye mixture according to claim 71, wherein each of k and l is zero, t is 1 and R^{61} is a phenyl group substituted by at least one sulphonic acid group or a salt thereof.

73. A dye mixture according to claim 71, wherein each of k, l and t is 1, each of R^{60} and R^{61} independently is a phenyl group

substituted by at least one sulphonic acid group or a salt thereof and Het² is a triazine ring substituted by a halogen atom.

74. A dye mixture according to claim 71, wherein each of k and t is zero, l is 1 and Het² is a reactive group having an aliphatic chain.

75. A dye mixture according to any one of claims 1 to 7, wherein the reactive dye (B) is a disazo dye of the formula (VIII), given and defined in claim 1.

76. A dye mixture according to claim 75, wherein

R⁵⁵ is a naphthyl group;

R⁵⁶ is a sulphonic acid group or a salt thereof;


R⁵⁹ is a group L¹⁰-Het, where L¹⁰ is the group N(H)-

and Het is a reactive heterocyclic group substituted by at least one halogen atom;

y₁ is 1, 2 or 3; and

each of y₂, y₃ and y₄ is zero.

77. A dye of the formula



78. A dye mixture according to claim 75, wherein

R^{55} is a phenyl group;

the group R^{56} or each group R^{56} independently is a sulphonic acid group or a salt thereof or is a reactive group selected from a vinyl sulphonyl group and a group $SO_2CH_2CH_2Q^2$ in which Q^2 is as defined in claim 1;


y_1 is 1 or 2; and,

at least one group R^{56} is a said reactive group or the group R^{59} is or includes a reactive group.

79. A dye mixture according to claim 78, wherein at least one group R^{56} or the group R^{59} is a reactive group selected from a vinyl sulphonyl group and a group $SO_2CH_2CH_2Q^2$ in which Q^2 is as defined in claim 1.


80. A dye mixture according to claim 78, wherein at least one group R^{56} is a sulphonic acid group or a salt thereof and R^{59} is the group L^{10} -Het, where L^{10} is the group $N(H)$ - and Het is a reactive triazine group substituted by a halogen atom and additionally, by the group $NHCH_2CH_2SO_3H$.

81. A disazo dye of the formula



(85A)

82. A disazo dye of the formula



83. A dye mixture according to any one of claims 1 to 7, wherein the reactive dye (B) is a phthalocyanine dye of the formula (X), given and defined in claim 1.

84. A dye mixture according to claim 83, wherein, in the reactive dye (B) of the formula (X), respective average values of x_1 , x_2 and x_3 are x_1 is 3, x_2 is zero and x_3 is 1, three of the four isoindole rings of the phthalocyanine have a respective sulphonic acid substituent (or a salt thereof) thereon and the other isoindole ring has a group $\text{SO}_2\text{NH-B-NH-Het}$ substituted thereon.

85. A dye mixture according to claim 84, wherein B is a straight or branched C_{2-4} alkylene group and Het is a triazine ring substituted by at least one halogen atom and optionally additionally substituted by a methoxy group.

86. A dye mixture according to any one of claims 1 to 7, wherein the reactive dye (B) is a triphendioxazine dye of the

formula (XI), given and defined in claim 1, or a salt thereof.

87. A dye mixture according to claim 86, wherein, in the reactive dye (B) of the formula (XI), each of T^1 and T^2 is a halogen atom, each U^1 is a sulphonic acid group or a salt thereof, each of B^1 and B^2 independently is a straight or branched C_{2-4} alkylene group, each of Het^1 and Het^2 independently is a triazine ring substituted by a halogen atom and each of R^{80} and R^{81} independently is a phenyl group substituted by at least one sulphonic acid group or a salt thereof.

88. A dye mixture according to any preceding claim comprising, by weight of the total weight of the dyes, from 10 to 99.5% by weight, inclusive of component (A), given and defined above, and from 0.5 to 90% by weight, inclusive of component (B), given and defined in claim 1.